



King's Research Portal

DOI:

[10.1093/cvr/cvt129](https://doi.org/10.1093/cvr/cvt129)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Kizub, I. V., Strielkov, I. V., Shaifta, Y., Becker, S., Prieto-Lloret, J., Snetkov, V. A., Soloviev, A. I., Aaronson, P. I., & Ward, J. P. T. (2013). Gap junctions support the sustained phase of hypoxic pulmonary vasoconstriction by facilitating calcium sensitization. *Cardiovascular Research*, 99(3), 404-411. <https://doi.org/10.1093/cvr/cvt129>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Gap junctions support the sustained phase of hypoxic pulmonary vasoconstriction by facilitating calcium sensitization

Igor V. Kizub^{1,2}, Ievgen V. Strielkov¹, Yasin Shaifta², Silke Becker², Jesus Prieto-Lloret², Vladimir A. Snetkov², Anatoly I. Soloviev¹, Philip I. Aaronson², and Jeremy P.T. Ward.²

¹Department of Experimental Therapeutics, Institute of Pharmacology and Toxicology of National Academy of Medical Sciences of Ukraine, Kiev, Ukraine; ²Division of Asthma, Allergy, and Lung Biology, King's College London, London, UK.

Corresponding author:

Prof Jeremy PT Ward

Division of Asthma, Allergy, and Lung Biology,

King's College London,

5th Floor Tower Wing, Guy's Campus

London SE1 9RT

Email: jeremy.ward@kcl.ac.uk

Tel/Fax: 020 7848 6695

© The Author 2013. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Abstract

Aims. To determine the role of gap junctions (GJ) in hypoxic pulmonary vasoconstriction (HPV).

Methods and results. Studies were performed in rat isolated intrapulmonary arteries (IPA) mounted on a myograph, and in anesthetized rats. Hypoxia induced a biphasic HPV response in IPA precontracted with prostaglandin $F_{2\alpha}$ ($PGF_{2\alpha}$, 3 μ M) or 20 mM K^+ . The GJ inhibitors 18 β -glycyrrhetic acid (18 β -GA, 30 μ M), heptanol (3.5 mM) or 2-aminoethoxydiphenyl borate (2-APB) (75 μ M) had little effect on the transient phase 1 of HPV, but abolished the sustained phase 2 which is associated with Ca^{2+} sensitisation. The voltage-dependent Ca^{2+} channel blocker diltiazem (10 μ M) had no effect on HPV, and did not alter the inhibitory action of 18 β -GA. Sustained HPV is enhanced by high glucose (15 mM) via potentiation of Ca^{2+} sensitization; in the presence of high glucose 18 β -GA still abolished sustained HPV. Simultaneous measurement of tension and intracellular Ca^{2+} using Fura PE-3 demonstrated that whilst 18 β -GA abolished tension development during sustained HPV, it did not affect the elevation of intracellular Ca^{2+} . Consistent with this, 18 β -GA abolished hypoxia-induced phosphorylation of the Rho kinase target MYPT-1. In anaesthetized rats hypoxia caused a biphasic increase in systolic right ventricular pressure. Treatment with oral 18 β -GA (25 mg/kg) abolished the sustained component of the hypoxic pressor response.

Conclusion. These results imply that GJs are critically involved in the signalling pathways leading to Rho kinase-dependent Ca^{2+} sensitization during sustained HPV, but not elevation of intracellular Ca^{2+} , and may explain the dependence of the former on an intact endothelium.

Key words: hypoxic pulmonary vasoconstriction; pulmonary artery; Ca^{2+} sensitization; gap junctions; 18 β -glycyrrhetic acid

Introduction

Hypoxic pulmonary vasoconstriction (HPV) optimizes pulmonary ventilation/perfusion matching in response to alveolar hypoxia, thereby diverting blood away from poorly ventilated regions of the lung. Despite many years of extensive research, the precise mechanisms underlying HPV remain incompletely resolved ¹.

In isolated intrapulmonary arteries (IPA), and some perfused lung preparations, the contractile response to acute hypoxia is typically biphasic, with a rapid transient vasoconstriction (phase 1) superimposed on a more slowly developing sustained contraction (phase 2) ^{1,2}Dipp, 2001 #2779;Robertson, 2003 #3055}. The mechanisms of these two phases differs. Phase 1 is associated with a transient elevation in intracellular Ca^{2+} ($[\text{Ca}^{2+}]_i$) involving store operated Ca^{2+} entry (SOCE) and Ca^{2+} influx through L-type voltage-dependent Ca^{2+} channels (VDCC), whereas the sustained phase 2 is critically dependent on ryanodine-sensitive Ca^{2+} stores and RhoA/Rho-kinase-dependent Ca^{2+} sensitization ^{1,3-6}. Intriguingly, both phase 2 of HPV and the associated Ca^{2+} sensitization have an yet unexplained strong dependence on an intact endothelium and glycolysis ^{2,4,7-9}.

Gap-junction-mediated communication within and between the endothelium and smooth muscle cells (SMCs) is important for the control and coordination of normal vascular function. Gap junctions (GJs) are formed of adjacent connexons between cells each comprised of 6 connexins (Cxs); the complete channel allows passage of electrical current and small signaling molecules ¹⁰⁻¹². Vascular tissues including pulmonary artery express Cx37, Cx40, Cx43 and Cx45 ^{11,13}. The endothelium couples to smooth muscle via myoendothelial GJs (MEGJs) ^{10,11}, which are present in IPA and probably play an important role in the integration of smooth muscle and endothelial function ¹³⁻¹⁵.

Inhibition of GJs suppresses HPV in perfused lungs ¹⁶, and it has recently been proposed that the signal for HPV originates at the alveolocapillary level, from which it is

propagated through the endothelium to upstream arterioles in a Cx40-dependent manner¹⁷. We have also reported that inhibition of glycolysis abolishes the hypoxia-induced depolarisation of pulmonary artery endothelial cells⁹, suggesting that GJs might be involved in the transduction pathways underlying the glycolysis and endothelium dependent sustained phase 2 of HPV.

The aim of the present study was to test the hypothesis that GJs contribute to local signalling in HPV, in addition to any upstream signal propagation, and to examine their role in the development of the glycolysis- and endothelium-dependent sustained phase of HPV.

2. Methods

2.1 Animals and tissue isolation

This study conforms with UK Home Office regulations and Directive 2010/63/EU of the European Parliament. Ethical approval was obtained from the relevant committees at King's College London and Institute of Pharmacology and Toxicology of National Academy of Medical Sciences of Ukraine. For isolated tissue studies, adult male Wistar rats (225–275g) were killed by lethal overdose of pentobarbital (i.p.) and cervical dislocation. The heart and lungs were excised and placed into cold physiological saline solution (PSS; in mM: 118 NaCl, 24 NaHCO₃, 1 MgSO₄, 0.44 NaH₂PO₄, 4 KCl, 5.5 glucose, and 1.8 CaCl₂).

2.2 Tension and intracellular Ca²⁺ measurements

Small IPA (150–400 µm in diameter) were dissected free of connective tissue, mounted on a wire myograph (Danish Myo Technology A/S, Aarhus, Denmark) and bathed in PSS gassed with 5% CO₂, balance air (pH 7.4) at 37°C. Vessels were stretched and pre-conditioned by

stimulation with repeated 2-min exposures to 80 mM K⁺ PSS (KPSS, equimolar substitution for NaCl) as previously described^{2,6,7}. Simultaneous measurement of IPA tension and intracellular Ca²⁺ ([Ca²⁺]_i) was performed using IPA mounted on a modified myograph, and following loading with Fura PE-3-AM (4 μM) for 1 h at 37°C. The myograph was placed on an inverted stage fluorescence microscope (Zeiss UK Ltd) with Fluor objective combined with a double-excitation microfluorimeter (CairnResearch Ltd, Faversham, UK). Tension was recorded simultaneously with light emitted by the artery at 510 nm at excitation wavelengths 340 and 380 nm. The ratio of emission intensities (R_{340/380}) was taken as a measure of smooth muscle [Ca²⁺]_i as described previously⁶⁻⁸. Tension and R_{340/380} were recorded using Acquisition Engine software (Cairn Research Ltd., UK).

2.3 Experimental protocols for in vitro studies

As previously described⁶⁻⁸, IPAs were pre-constricted with sufficient PGF_{2α} to produce tension equivalent to 10–15% of that produced by KPSS (typically 3 μM), in order to elicit a full contractile response to hypoxia. In some experiments equivalent pretone was induced with PSS containing 20–25 mM [K⁺]. Hypoxia was induced by switching from 95% air/5% CO₂ to 5% CO₂/balance N₂, which we have shown to provide a pO₂ of 15–20 mmHg during hypoxia and 145–150 mmHg during normoxia in the myograph chamber³. Consecutive hypoxic challenges, separated by 60 min normoxia, demonstrated highly consistent HPV responses within the same IPA. Tension and changes in [Ca²⁺]_i (R_{340/380}) are expressed as percentages of the maximal responses to contraction to KPSS.

IPA incubated under the same conditions as above were snap frozen and protein extracted for western blot analysis of myosin phosphatase targeting protein (MYPT-1) phosphorylation as previously described⁶. Membranes were blocked for 1h with 5% milk, probed with primary antibody overnight at 4°C (1:1000–1:5000 in 5% milk), followed by application of

horseradish peroxidase-conjugated anti-IgG secondary antibody (Sigma, UK) for 1h at room temperature (1:5000). Membranes were first probed with anti-phospho-MYPT-1 (thr-855; Upstate, UK), stripped for 1h (Pierce stripping buffer), re-blocked, and re-probed with anti-pan MYPT-1 (Cell Signalling, UK). Data are expressed as ratio phospho/pan MYPT-1 as a percentage of in-gel controls.

2.4 In vivo studies

Experiments were conducted on adult male Wistar rats (230–300g) divided into control and treatment groups. Plasma concentrations of 18β -GA in rats have been shown to fall rapidly after oral administration, but after ~12 hrs become relatively stable for up to 24 hr¹⁸. Animals were therefore treated orally with 18β -GA (25 mg/kg) 20 hours before experimentation. Surgical anaesthesia was induced by intra-peritoneal injection of chloralose-urethane (1:10; 40 mg of urethane per 100 g body weight). Once deep anaesthesia was confirmed, tracheal intubation was performed. The left jugular vein and left common carotid artery were catheterized, and heparin (50 U per 100 g body weight) infused. Catheterization of the right ventricle was performed through the right jugular vein. Right ventricular and carotid artery pressures were recorded with ISOTEC pressure transducers (HSE, Germany) and Chart 5 Pro (ADInstruments Ltd, Australia). Animals were mechanically ventilated with a minute volume of 140 ml/min (Ugo Basile 7025 ventilator), and initial values of parameters recorded for ~25 min after stabilization. Hypoxia was then induced for 30 min by ventilation with 10% O₂ in N₂. Animals were euthanized at the end of the experiment by means of intravenous urethane (400 mg/100 g).

2.5 Statistical analysis

Results are expressed as means \pm SEM. Statistical analysis was performed using ANOVA with a Holm-Sidak post hoc test or Student's t test as appropriate (Sigmaplot 12, Systat Software Inc., CA).

2.6. Chemicals

Diltiazem, carbachol, 18β -glycyrrhetic acid (18β -GA), heptanol, 2-aminoethoxydiphenyl borate (2-APB) and Fura-PE-3/AM were obtained from Sigma-Aldrich UK. All drugs were dissolved in deionized water except 18β -GA, heptanol and Fura PE-3 AM, which were dissolved in DMSO to make stock solutions; following final dilution DMSO had no effect on its own.

3. Results.

3.1 Effect of gap junction inhibitors on HPV in isolated IPA

We examined the effects of 3 structurally unrelated GJ inhibitors which are thought to act by different mechanisms, 18β -glycyrrhetic acid (18β -GA), heptanol, and 2-aminoethoxydiphenyl borate (2-APB)^{19,20}; whilst 2-APB is commonly used as an inhibitor of SOCE^{21,22}, it is also an effective blocker of GJ, with differential efficacy according to the Cx of which they are formed¹⁹. The synthetic^{37,43} Gap27 GJ inhibitor had inconsistent effects, possibly due to diffusional limitations, and was not pursued.

As previously described, hypoxia elicited a biphasic response in tension in IPA precontracted with $\text{PGF}_{2\alpha}$ (Fig. 1). The first phase consisted of a transient contraction, which peaked within 3–5 min of the onset of hypoxia, followed by a relaxation which reached a nadir within 10–15 min. This was followed by a more slowly developing and sustained contraction (phase 2). Reoxygenation resulted in a rapid return of tension to the initial $\text{PGF}_{2\alpha}$ -induced values, and on washing with PSS tension returned to the original baseline.

Preincubation with 18 β -GA (30 μ M) was without effect on phase 1 of HPV, but strongly suppressed phase 2 ($p < 0.001$; Fig.1). The effect of 18 β -GA on HPV was independent of the means of precontraction, as equivalent results were obtained in IPA precontracted by depolarization with PSS containing 20 mM [K⁺] (Fig. 1). Both heptanol (3.5 mM) and 2-APB (75 μ M) also strongly suppressed phase 2; although both caused a small reduction in phase 1, this only reached significance for heptanol (Fig. 2). A lower concentration of 2-APB (30 μ M) reduced phase 2 at 40 min by only 58 \pm 6% ($n = 5$, $p < 0.01$).

Together these data suggest that GJ are involved in the sustained phase 2 of HPV, but not the transient phase 1.

3.2 Effect of 18 β -glycyrrhetic acid on HPV following blockade of L-type Ca²⁺ channels

Blockade of GJs could potentially affect membrane potential in the smooth muscle. We therefore compared the control HPV response to that following incubation with the L-type VDCC blocker diltiazem (10 μ M), and in combination with 18 β -GA (30 μ M) (Fig. 3). As previously reported³, diltiazem alone had no significant effect on either phase of HPV in this preparation; the reduction in phase 1 did not reach significance (control: 28.7 \pm 2.6% KPSS; diltiazem: 23.4 \pm 2.3% KPSS, $n = 7$, NS). However, addition of 18 β -GA to diltiazem strongly suppressed the sustained phase 2 of HPV (Fig. 3). Whilst phase 1 was now significantly reduced compared to control ($p < 0.05$), it was not significantly different from that with diltiazem alone (diltiazem + 18 β -GA: 18.8 \pm 2.7% KPSS, $n = 7$, NS). These results, which are essentially equivalent to those shown in Figure 1, demonstrate that the contribution of GJs to sustained HPV is independent of VDCC.

3.3 Effect of 18 β -glycyrrhetic acid on intracellular Ca²⁺ concentration during HPV.

Hypoxia elicited a biphasic response in $[Ca^{2+}]_i$ in IPA precontracted with 3 μ M $PGF_{2\alpha}$, with a transient increase in $[Ca^{2+}]_i$ that mirrored the phase 1 transient increase in tension (Fig. 4). However, as previously reported^{7,8,23}, whereas tension gradually increased during phase 2, $[Ca^{2+}]_i$ remained stable. Reoxygenation resulted in a rapid return of both tension and $[Ca^{2+}]_i$ to the initial $PGF_{2\alpha}$ -induced values. Application of 18 β -GA (30 μ M) had no effect on hypoxia-induced elevation of $[Ca^{2+}]_i$ during either phase of HPV (Fig. 4). This suggests that GJs are involved in the mechanisms leading to Ca^{2+} sensitization during sustained HPV, but not the elevation of $[Ca^{2+}]_i$.

3.4 Effect of 18 β -glycyrrhetic acid on HPV in the presence of elevated glucose

We have previously shown that phase 2 of HPV is selectively potentiated by an increase in extracellular glucose concentration, and suppressed or abolished by reduced glucose; this involves the mechanisms underlying the phase 2 associated Ca^{2+} sensitization, as altering glucose had no effect on the hypoxia-induced elevation of $[Ca^{2+}]_i$ ⁷. As our results suggest that GJs are also involved in Ca^{2+} sensitization during phase 2, we examined the effects of GJ inhibition on the high glucose-induced potentiation of phase 2 of HPV. Increasing glucose concentration from 5 to 15 mM had no significant effect on basal tension, pretone elicited by $PGF_{2\alpha}$, or HPV phase 1, but phase 2 was significantly potentiated (Fig. 5A). In the presence of 15 mM glucose, 18 β -GA (30 μ M) again had no effect on phase 1 but abolished phase 2 (Fig. 5).

This suggests that the mechanisms underlying the potentiating action of high glucose on phase 2 of HPV are not separate from those inhibited by block of GJs, and strengthens the hypothesis that Ca^{2+} sensitisation during hypoxia requires functional GJs.

3.4 Effect of 18 β -glycyrrhetic acid on MYPT-1 phosphorylation during HPV.

Ca^{2+} sensitization during sustained HPV depends on Rho kinase-dependent phosphorylation of MYPT-1^{6,24}. We therefore examined the effect of 18 β -GA on MYPT-1 phosphorylation in IPA precontracted with PGF_{2 α} as above and following 30 min hypoxia. In controls, hypoxia increased the ratio of phosphorylated to total MYPT-1 by ~30% compared to PGF_{2 α} alone ($p < 0.05$, $n = 10$). This hypoxia-induced increase was however abolished following treatment with 18 β -GA, although the response to precontraction with PGF_{2 α} alone was unaltered by 18 β -GA ($n = 8-9$; Fig. 5B). These results further suggest that blockade of GJs with 18 β -GA impairs hypoxia-induced Rho kinase-mediated Ca^{2+} sensitisation.

3.5 Effect of 18 β -glycyrrhetic acid on right ventricular pressure *in vivo*.

Basal systolic right ventricular pressure (sRVP) in untreated anaesthetized rats was 28.6 ± 1.1 mmHg and mean systemic arterial pressure (MAP) 101.5 ± 5.3 mm Hg ($n = 7$). Induction of hypoxia resulted in a biphasic pressor response (Fig. 6). sRVP rose to a peak of ~44% (12.6 ± 0.8 mmHg, $p < 0.001$) after 5 ± 1 min, then fell to a nadir of ~24% (7.2 ± 4.2 mmHg, $p < 0.001$ vs peak) before increasing again to ~38 % by 32 min (10.9 ± 0.8 mmHg, $p < 0.001$ vs baseline and nadir) (Fig. 6). Hypoxia caused a fall in MAP over the first 3 min to 64.8 ± 5.4 mmHg ($p < 0.001$), which thereafter remained constant throughout the period of hypoxic. All changes in haemodynamics reversed on re-oxygenation after ~5-10 minutes.

Following oral administration of 18 β -GA (25 mg/kg) 20 hours before the study, basal sRVP was reduced compared to controls, though this did not reach significance (26.0 ± 1.4 mmHg, $n = 5$, NS). Conversely there was significant elevation in MAP (120.5 ± 4.5 mmHg, $p < 0.05$). The initial elevation of sRVP following induction of hypoxia was reduced to 6.9 ± 4.3 mmHg, though this did not reach significance; however the sustained component of the

hypoxic pressor response was effectively abolished (32 min: 0.9 ± 1.9 mmHg, $p < 0.001$) (Fig. 6). MAP during hypoxia showed no difference between control and treated animals (control: 64.8 ± 5.4 mm Hg; 18β -GA: 67.0 ± 8.2 mmHg, NS).

4. Discussion

The key finding of this study is that the sustained phase 2 of HPV, but not the transient phase 1, is critically dependent on functioning GJ, and our evidence suggests a hitherto unrecognized role for GJs in hypoxia-induced Ca^{2+} sensitization. As this dependence was apparent in isolated IAP segments, it must be localized to the artery wall and independent of any longitudinal signal propagation from downstream regions of the pulmonary vascular tree.

The biphasic nature of HPV has been recognized for many years, and the initial phase 1 constriction differs mechanistically from that during the sustained phase 2¹. Although both phases are associated with an elevation of smooth muscle $[\text{Ca}^{2+}]_i$, this is very small in phase 2 and critically dependent on release from ryanodine-sensitive stores, with a reduced or absent dependence on Ca^{2+} entry^{3,4,25}. In particular, and in contrast to phase 1, tension development during phase 2 is strongly dependent on the endothelium and Rho kinase-mediated Ca^{2+} sensitization^{4-6,8,23}.

Whilst this biphasic response is most clearly demonstrated in isolated small IPA^{2,4,8,26}, it is occasionally apparent in perfused lung preparations and *in vivo* where the hypoxic challenge is prolonged for more than 20 min^{1,27,28}. However, the majority of studies on HPV, in whatever preparation, utilize shorter hypoxic challenges of up to 15 min, and are thus effectively only relevant to phase 1^{1,25}. Whilst the relative importance of the two phases for acute ventilation-perfusion matching is a subject of debate, the sustained phase 2 is more relevant to group 3 pulmonary hypertension with which it shares some common features, specifically a key role for Rho kinase-mediated Ca^{2+} sensitization²⁴.

As our studies were designed to investigate a role for GJs in local signalling in the absence of signal propagation from remote regions of the pulmonary vasculature, the key experiments were methodologically constrained to small, endothelium-intact segments of

IPA. This prevented a molecular approach as incubation of such preparations for sufficient time to allow transfection with e.g. siRNAs causes significant changes in function (e.g.²⁹ and own observations). It is also notoriously difficult to induce HPV in isolated mouse IPA, obviating use of GM models¹. We therefore utilized a pharmacological approach with three GJ inhibitors that are believed to act via different mechanisms, 18 β -GA, heptanol and 2-APB. All three demonstrated a similar profile, with strong suppression of HPV phase 2 and little or no effect on phase 1 (Fig. 1 and 2). Notably, the hypoxic pressor response *in vivo* also showed a biphasic response, and treatment with 18 β -GA similarly abolished the sustained component of HPV but not the initial transient (Fig. 6).

18 β -GA and heptanol are widely used as GJ inhibitors in vascular tissues^{19,30-32}. 18 β -GA is a triterpenoid saponin that disrupts gap junction plaques by affecting the structural integrity, distribution and phosphorylation of connexons^{19,32}. The long-chain alcohol heptanol causes selective disruption at the GJ-lipid interface by intercalating in the lipid bilayer and thus effectively gating GJ channels closed^{19,30-32}. The mechanism of 2-APB is less clear, but it is recognized as a potent blocker of GJs with a much greater efficacy against Cx40 compared to Cx43^{19,20}. It is commonly used as an inhibitor of IP₃ receptors (which do not play a major role in HPV¹), and SOCE^{21,22}, although notably it is reported that 75 μ M 2-APB only inhibits SOCE in SMCs isolated from IPA by ~40%³³.

Whilst SOCE has been implicated in HPV, the evidence is largely restricted to phase 1 and studies in isolated PASMCs^{1,3,34}. SOCE is highly sensitive to La³⁺, and in IPA 1 μ M La³⁺ abolished thapsigargin-induced SOCE and suppressed phase 1 of HPV^{3,22}; 10 μ M La³⁺ also suppressed the acute hypoxic pressor response in perfused lungs³⁴. However during phase 2 of HPV in IPA, 1 μ M La³⁺ was without effect on the elevation of either tension or [Ca²⁺]_i, and La³⁺ only caused partial suppression at 100 μ M, suggesting that SOCE plays a limited role during this phase³. Notably, at concentrations of 100 μ M and above La³⁺ blocks

Cx43-containing GJs³⁵. In the light of this, and as the effect of 2-APB on HPV showed a similar profile to that of 18 β -GA and heptanol with a strong and selective block of phase 2, it is reasonable to suggest that its actions here are mediated by its established ability to block GJs^{19,20}.

Inhibitors of GJs, including 18 β -GA and Cx mimetic inhibitory peptides (Gap27), have been reported to elevate systemic blood pressure *in vivo*, as observed here. This has been ascribed to inhibition of EDHF-dependent vasorelaxation and/or interference with the renin-angiotensin system^{31,36}. In contrast GJ inhibitors *in vitro* generally induce vasorelaxation or attenuation of constriction, due to smooth muscle hyperpolarization, inhibition of Ca²⁺ entry, and a consequent fall in [Ca²⁺]_i^{31,32,37}. As HPV has been associated with hypoxia-induced Ca²⁺ entry^{1,3,27}, it is feasible that GJ inhibition could be suppressing phase 2 in a similar fashion. However, as previously reported³ block of voltage gated L-type Ca²⁺ channels had no effect on either phase of HPV in this preparation, and 18 β -GA still selectively abolished phase 2 (Fig.5). Moreover, whilst 18 β -GA strongly inhibited tension development during phase 2, it did not affect the hypoxia-induced elevation of [Ca²⁺]_i (Fig. 4). This implies that inhibition of GJs is suppressing hypoxia-induced Ca²⁺ sensitization.

Phase 2 of HPV involves Rho kinase-mediated Ca²⁺ sensitization, which is endothelium- and glucose-dependent^{1,5,7,8}. Endothelial denudation or removal of glucose abolishes phase 2 tension development, whilst high glucose potentiates it, in all cases without affecting the associated elevation of [Ca²⁺]_i; notably, high glucose cannot restore HPV in the absence of endothelium⁵⁻⁸. If functioning GJs are required for hypoxia-induced Ca²⁺ sensitization, it would be predicted that high glucose would have no effect following GJ blockade with 18 β -GA, and this is what was observed (Fig. 5A). Moreover 18 β -GA abolished hypoxia-induced phosphorylation of MYPT-1, the target for Rho kinase and a key regulator of Ca²⁺ sensitivity (Fig. 5B).

Our results are consistent with a hitherto unrecognised role for GJs in mediating hypoxia-induced and Rho kinase dependent Ca^{2+} sensitisation during HPV, which could underlie the so far unexplained dependence of phase 2 in isolated arteries on an intact endothelium. It has been hypothesised that hypoxia causes release of an endothelium derived constricting factor (EDCF), but no such factor with the requisite characteristics has been positively identified; whilst endothelin 1 and others may play a role in HPV, possibly by providing the endogenous equivalent of pretone, they have been excluded as the mediator of phase 2 Ca^{2+} sensitisation^{1,38,39}. MEGJ allow direct passage of small signalling molecules up to ~1.2 kDa¹². Intriguingly, Gairhe et al. have recently shown that serotonin synthesised in pulmonary vascular endothelial cells passes through MEGJs to modify smooth muscle differentiation⁴⁰, and binding of intracellular serotonin to RhoA (serotonylation) has been associated with the increased smooth muscle RhoA/Rho kinase activity observed in pulmonary hypertension⁴¹.

Blockade of MEGJs has been reported to inhibit contraction of IPA to exogenous serotonin and other agonists by preventing transfer of reactive oxygen species (ROS), most likely superoxide anion¹³. This is interesting because of strong evidence that ROS comprise the key mediator of HPV^{1,24,42}, and activate Rho kinase^{43,44}. Diacylglycerol (DAG) is another small signalling molecule implicated in HPV. Although interest has focussed on its activation of TRPC6 channels during the acute phase^{27,45}, inhibition of phosphatidylcholine-specific phospholipase C, potentially the source of DAG during hypoxia, is reported to selectively suppress sustained HPV^{26,46}.

GJs between endothelial cells allow longitudinal propagation of electrical signals along small vessels. A recent study by Wang et al. suggests that retrograde propagation via Cx40-containing GJs between endothelial cells couples oxygen sensing in alveolar capillaries

to contraction of upstream pulmonary arterioles, and may thus be a critical requirement for HPV¹⁷. Such a mechanism is clearly not applicable to the short segments of IPA used in the current study. It also seems unlikely that Cx40 is critically involved in the responses we describe, as unlike Cx43 and most others, Cx40 is effectively completely blocked by 20 μ M 2-APB²⁰, whereas in our hands 30 μ M 2-APB only caused partial inhibition of phase 2. In any event, electrotonic coupling is unlikely to play a role in HPV of isolated IPA, as we have previously demonstrated that this is essentially unaltered following near-maximal depolarisation with 80mM $[K^+]$ ³. However, our studies do not rule out a role for longitudinal propagation in HPV in the intact lung, as proposed by Wang et al.¹⁷.

In conclusion, our results suggest that GJs, probably MEGJs, play a critical and previously unsuspected role in sustained HPV by facilitating hypoxia-induced Ca^{2+} sensitization. This is independent of any role in longitudinal signal propagation as it is apparent in isolated IPA segments. Whilst the GJ inhibitors utilised also have non-junctional actions^{20,37}, their disparate *modi operandi* yet identical actions on HPV, with little effect on phase 1 but robust inhibition of phase 2, strongly suggests that the latter is indeed due to block of GJs. The mechanism however remains unclear, but is likely to involve transfer of a small signalling molecule such as serotonin or superoxide between endothelium and smooth muscle rather than electrical coupling.

Funding

This work was supported by a Physiological Society Junior Fellowship Grant (IVK); Royal Society International Travel Grant for Collaboration (IVK); and the Wellcome Trust (grant #087776, JPTW & PIA).

Conflict of Interest: None declared

References

1. Sylvester JT, Shimoda LA, Aaronson PI, Ward JP. Hypoxic pulmonary vasoconstriction. *Physiol Rev* 2012;**92**:367-520.
2. Leach RM, Robertson TP, Twort CH, Ward JP. Hypoxic vasoconstriction in rat pulmonary and mesenteric arteries. *Am J Physiol Lung Cell Mol Physiol* 1994;**266**:L223-231.
3. Robertson TP, Hague D, Aaronson PI, Ward JP. Voltage-independent calcium entry in hypoxic pulmonary vasoconstriction of intrapulmonary arteries of the rat. *J Physiol* 2000;**525**:669-680.
4. Dipp M, Nye PC, Evans AM. Hypoxic release of calcium from the sarcoplasmic reticulum of pulmonary artery smooth muscle. *Am J Physiol Lung Cell Mol Physiol* 2001;**281**:L318-325.
5. Robertson TP, Dipp M, Ward JP, Aaronson PI, Evans AM. Inhibition of sustained hypoxic vasoconstriction by Y-27632 in isolated intrapulmonary arteries and perfused lung of the rat. *Br J Pharmacol* 2000;**131**:5-9.
6. Knock GA, Snetkov VA, Shaifta Y, Drndarski S, Ward JP, Aaronson PI. Role of src-family kinases in hypoxic vasoconstriction of rat pulmonary artery. *Cardiovasc Res* 2008;**80**:453-462.
7. Leach RM, Hill HM, Snetkov VA, Robertson TP, Ward JP. Divergent roles of glycolysis and the mitochondrial electron transport chain in hypoxic pulmonary vasoconstriction of the rat: identity of the hypoxic sensor. *J Physiol* 2001;**536**:211-224.
8. Robertson TP, Aaronson PI, Ward JP. Ca²⁺ sensitization during sustained hypoxic pulmonary vasoconstriction is endothelium dependent. *Am J Physiol Lung Cell Mol Physiol* 2003;**284**:L1121-1126.
9. Soloviev AI, Bondarenko AI, Kizub IV. Selective glycolysis blockade in guinea pig pulmonary artery and aorta reverses contractile and electrical responses to acute hypoxia. *Vascul Pharmacol* 2012;**57**:119-123.
10. Figueroa XF, Duling BR. Gap junctions in the control of vascular function. *Antioxidants & redox signaling* 2009;**11**:251-266.
11. Brisset AC, Isakson BE, Kwak BR. Connexins in vascular physiology and pathology. *Antioxidants & redox signaling* 2009;**11**:267-282.
12. Saez JC, Berthoud VM, Branes MC, Martinez AD, Beyer EC. Plasma membrane channels formed by connexins: their regulation and functions. *Physiol Rev* 2003;**83**:1359-1400.
13. Billaud M, Marthan R, Savineau JP, Guibert C. Vascular smooth muscle modulates endothelial control of vasoreactivity via reactive oxygen species production through myoendothelial communications. *PloS one* 2009;**4**:e6432.

14. Davies P, Burke G, Reid L. The structure of the wall of the rat intraacinar pulmonary artery: an electron microscopic study of microdissected preparations. *Microvasc Res* 1986;**32**:50-63.
15. Billaud M, Dahan D, Marthan R, Savineau JP, Guibert C. Role of the gap junctions in the contractile response to agonists in pulmonary artery from two rat models of pulmonary hypertension. *Respiratory Research* 2011;**12**:30.
16. Morio Y, Carter EP, Oka M, McMurtry IF. EDHF-mediated vasodilation involves different mechanisms in normotensive and hypertensive rat lungs. *Am J Physiol Heart Circ Physiol* 2003;**284**:H1762-1770.
17. Wang L, Yin J, Nickles HT, Ranke H, Tabuchi A, Hoffmann J, *et al.* Hypoxic pulmonary vasoconstriction requires connexin 40-mediated endothelial signal conduction. *J Clin Invest* 2012;**122**:4218-4230.
18. Sun HY, Li Q, Chen W, Geng LL, Li X, Chen XH, *et al.* Pharmacokinetic analysis of alpha and beta epimers of glycyrrhetic acid in rat plasma: differences in singly and combined administrations. *Yao Xue Xue Bao* 2012;**47**:94-100.
19. Dhein S. Pharmacology of gap junctions in the cardiovascular system. *Cardiovasc Res* 2004;**62**:287-298.
20. Bai D, del Corso C, Srinivas M, Spray DC. Block of specific gap junction channel subtypes by 2-aminoethoxydiphenyl borate (2-APB). *J Pharmacol Exp Ther* 2006;**319**:1452-1458.
21. Bootman MD, Collins TJ, Mackenzie L, Roderick HL, Berridge MJ, Peppiatt CM. 2-aminoethoxydiphenyl borate (2-APB) is a reliable blocker of store-operated Ca^{2+} entry but an inconsistent inhibitor of InsP_3 -induced Ca^{2+} release. *FASEB J* 2002;**16**:1145-1150.
22. Snetkov VA, Aaronson PI, Ward JP, Knock GA, Robertson TP. Capacitative calcium entry as a pulmonary specific vasoconstrictor mechanism in small muscular arteries of the rat. *Br J Pharmacol* 2003;**140**:97-106.
23. Robertson TP, Aaronson PI, Ward JP. Hypoxic vasoconstriction and intracellular Ca^{2+} in pulmonary arteries: evidence for PKC-independent Ca^{2+} sensitization. *Am J Physiol Heart Circ Physiol* 1995;**268**:H301-307.
24. Ward JP, McMurtry IF. Mechanisms of hypoxic pulmonary vasoconstriction and their roles in pulmonary hypertension: new findings for an old problem. *Curr Opin Pharmacol* 2009;**9**:287-296.
25. Aaronson PI, Robertson TP, Knock GA, Becker S, Lewis TH, Snetkov V, *et al.* Hypoxic pulmonary vasoconstriction: mechanisms and controversies. *J Physiol* 2006;**570**:53-58.
26. Strielkov IV, Khromov AS. Hypoxic pulmonary hypertension: The role of phosphatidylcholine-specific phospholipase C. *Int J Phys Pathophys* 2010;**1**:119-124.

27. Weissmann N, Dietrich A, Fuchs B, Kalwa H, Ay M, Dumitrascu R, *et al.* Classical transient receptor potential channel 6 (TRPC6) is essential for hypoxic pulmonary vasoconstriction and alveolar gas exchange. *Proc Natl Acad Sci U S A* 2006;**103**:19093-19098.
28. Talbot NP, Balanos GM, Dorrington KL, Robbins PA. Two temporal components within the human pulmonary vascular response to approximately 2 h of isocapnic hypoxia. *J Appl Physiol* 2005;**98**:1125-1139.
29. Manoury B, Etheridge SL, Reid J, Gurney AM. Organ culture mimics the effects of hypoxia on membrane potential, K(+) channels and vessel tone in pulmonary artery. *Br J Pharmacol* 2009;**158**:848-861.
30. Christ GJ. Modulation of alpha 1-adrenergic contractility in isolated vascular tissues by heptanol: a functional demonstration of the potential importance of intercellular communication to vascular response generation. *Life Sci* 1995;**56**:709-721.
31. Lagaud G, Davies KP, Venkateswarlu K, Christ GJ. The physiology, pathophysiology and therapeutic potential of gap junctions in smooth muscle. *Curr Drug Targets* 2002;**3**:427-440.
32. Earley S, Resta TC, Walker BR. Disruption of smooth muscle gap junctions attenuates myogenic vasoconstriction of mesenteric resistance arteries. *American Journal of Physiology - Heart and Circulatory Physiology* 2004;**287**:H2677-2686.
33. McElroy SP, Gurney AM, Drummond RM. Pharmacological profile of store-operated Ca(2+) entry in intrapulmonary artery smooth muscle cells. *Eur J Pharmacol* 2008;**584**:10-20.
34. Weigand L, Foxson J, Wang J, Shimoda LA, Sylvester JT. Inhibition of hypoxic pulmonary vasoconstriction by antagonists of store-operated Ca²⁺ and nonselective cation channels. *Am J Physiol Lung Cell Mol Physiol* 2005;**289**:L5-L13.
35. Contreras JE, Saez JC, Bukauskas FF, Bennett MV. Gating and regulation of connexin 43 (Cx43) hemichannels. *Proc Natl Acad Sci U S A* 2003;**100**:11388-11393.
36. Takenaka T, Inoue T, Kanno Y, Okada H, Meaney KR, Hill CE, *et al.* Expression and role of connexins in the rat renal vasculature. *Kidney Int* 2008;**73**:415-422.
37. Matchkov VV, Rahman A, Peng H, Nilsson H, Aalkjaer C. Junctional and nonjunctional effects of heptanol and glycyrrhetic acid derivatives in rat mesenteric small arteries. *Br J Pharmacol* 2004;**142**:961-972.
38. Gaine SP, Hales MA, Flavahan NA. Hypoxic pulmonary endothelial cells release a diffusible contractile factor distinct from endothelin. *Am J Physiol Lung Cell Mol Physiol* 1998;**274**:L657-664.
39. Robertson TP, Ward JP, Aaronson PI. Hypoxia induces the release of a pulmonary-selective, Ca²⁺-sensitising, vasoconstrictor from the perfused rat lung. *Cardiovasc Res* 2001;**50**:145-150.

40. Gairhe S, Bauer NN, Gebb SA, McMurtry IF. Serotonin passes through myoendothelial gap junctions to promote pulmonary arterial smooth muscle cell differentiation. *Am J Physiol Lung Cell Mol Physiol* 2012;**303**:L767-777.
41. Guilluy C, Eddahibi S, Agard C, Guignabert C, Izikki M, Tu L, *et al.* RhoA and Rho kinase activation in human pulmonary hypertension: role of 5-HT signaling. *Am J Respir Crit Care Med* 2009;**179**:1151-1158.
42. Waypa GB, Chandel NS, Schumacker PT. Model for hypoxic pulmonary vasoconstriction involving mitochondrial oxygen sensing. *Circ Res* 2001;**88**:1259-1266.
43. Knock GA, Snetkov VA, Shaifta Y, Connolly M, Drndarski S, Noah A, *et al.* Superoxide constricts rat pulmonary arteries via Rho-kinase-mediated Ca^{2+} sensitization. *Free Radic Biol Med* 2009;**46**:633-642.
44. Chi AY, Waypa GB, Mungai PT, Schumacker PT. Prolonged hypoxia increases ROS signaling and RhoA activation in pulmonary artery smooth muscle and endothelial cells. *Antioxidants & redox signaling* 2010;**12**:1-7.
45. Fuchs B, Rupp M, Ghofrani HA, Schermuly RT, Seeger W, Grimminger F, *et al.* Diacylglycerol regulates acute hypoxic pulmonary vasoconstriction via TRPC6. *Respiratory Research* 2011;**12**:20.
46. Strielkov IV, Ward JP, Aaronson PI. Evidence of phosphatidylcholine-specific phospholipase C involvement in hypoxic pulmonary vasoconstriction. *Proc Physiol Soc* 2011;**23**:PC358.

Figure legends

Figure 1: HPV in rat isolated IPA, and the effect of preincubation with 18β -GA ($30\ \mu\text{M}$) on the response. Panel A shows example traces following preconstriction with $3\ \mu\text{M}$ $\text{PGF}_{2\alpha}$, and the selective suppression of phase 2 by 18β -GA; mean data are shown in panel B ($n = 10$). 18β -GA had the same effect in IPA precontracted by depolarisation with $20\ \text{mM}$ $[\text{K}^+]$ (panel C, $n = 7$). Symbols represents mean \pm SE. * = $p < 0.05$, ** = $p < 0.01$.

Figure 2: The effect of the GJ inhibitors heptanol ($3.5\ \text{mM}$, panel A, $n = 6$) and 2-APB ($75\ \mu\text{M}$, panel B, $n = 7$) on HPV in rat IPA precontracted with $3\ \mu\text{M}$ $\text{PGF}_{2\alpha}$. Symbols represents mean \pm SE. * = $p < 0.05$, ** = $p < 0.001$.

Figure 3: Blockade of VDCC with diltiazem ($10\ \mu\text{M}$) had no significant effect on HPV in rat IPA precontracted with $3\ \mu\text{M}$ $\text{PGF}_{2\alpha}$ (filled circles); 18β -GA ($30\ \mu\text{M}$) had almost identical effects in the presence of diltiazem as in its absence (Fig. 1). Symbols represents mean \pm SE, $n = 7$. * = $p < 0.05$, ** = $p < 0.001$.

Figure 4: Simultaneous measurement of tension and $[\text{Ca}^{2+}]_i$ (expressed as $R_{340/380}$) during HPV in rat IPA precontracted with $3\ \mu\text{M}$ $\text{PGF}_{2\alpha}$, and following preincubation with 18β -GA ($30\ \mu\text{M}$). 18β -GA abolished the rise in tension phase 2, but had no effect on $[\text{Ca}^{2+}]_i$. Symbols represent mean \pm SE, $n = 7$. * = $p < 0.05$, ** = $p < 0.01$.

Figure 5: Panel A: Elevation of glucose concentration from 5 to $15\ \text{mM}$ potentiated phase 2 of HPV in rat IPA precontracted with $3\ \mu\text{M}$ $\text{PGF}_{2\alpha}$ (filled circles, $n = 7$, # = $p < 0.05$). In the presence of $15\ \text{mM}$ glucose 18β -GA still abolished phase 2 (filled squares, * = $p < 0.05$, ** =

$p < 0.01$, comparison with high glucose alone). Symbols represent mean \pm SE; where error bars are not shown, they are smaller than the symbol. Panel B: In the absence of 18β -GA, MYPT-1 phosphorylation (expressed as % unstimulated) in IPA was increased by precontraction with $\text{PGF}_{2\alpha}$ ($p < 0.05$) and further increased after 30 min hypoxia (* = $p < 0.05$ $\text{PGF}_{2\alpha}$ vs $\text{PGF}_{2\alpha}$ plus hypoxia; $n = 10$). In the presence of 18β -GA, the increase in MYPT-1 phosphorylation induced by precontraction with $\text{PGF}_{2\alpha}$ was unaltered, but thereafter hypoxia had no further effect (NS, $\text{PGF}_{2\alpha}$ vs $\text{PGF}_{2\alpha}$ plus hypoxia; $n = 8-9$).

Figure 6: Hypoxic pressor response in anaesthetized rats *in vivo*, and following oral administration of 25 mg/kg 18β -GA. Panel A shows example traces from a control and a treated animal, panel B mean data. In control animals hypoxia elicited a rapid elevation in sRVP followed by a small but significant decline, after which sRVP rose more slowly towards a sustained plateau (# = $p < 0.01$ between points as shown, $n = 7$). This sustained component was abolished in 18β -GA treated animals ($n = 5$). Symbols represent mean \pm SE, * = $p < 0.01$ (control vs treated). NS = not significant ($p > 0.05$).











